

RESEARCH

Open Access



Machine-learning-based models for the optimization of post-cervical spinal laminoplasty outpatient follow-up schedules

Yechan Seo^{1,2†}, Seoi Jeong^{2,3†}, Siyoung Lee⁴, Tae-Shin Kim⁵, Jun-Hoe Kim^{6,7}, Chun Kee Chung^{6,7,8}, Chang-Hyun Lee^{6,7}, John M. Rhee⁹, Hyoun-Joong Kong^{1,2,3*}  and Chi Heon Kim^{6,7,10*}

Abstract

Background Patients undergo regular clinical follow-up after laminoplasty for cervical myelopathy. However, those whose symptoms significantly improve and remain stable do not need to conform to a regular follow-up schedule. Based on the 1-year postoperative outcomes, we aimed to use a machine-learning (ML) algorithm to predict 2-year postoperative outcomes.

Methods We enrolled 80 patients who underwent cervical laminoplasty for cervical myelopathy. The patients' Japanese Orthopedic Association (JOA) scores (range: 0–17) were analyzed at the 1-, 3-, 6-, and 12-month postoperative timepoints to evaluate their ability to predict the 2-year postoperative outcomes. The patient acceptable symptom state (PASS) was defined as a JOA score ≥ 14.25 at 24 months postoperatively and, based on clinical outcomes recorded up to the 1-year postoperative timepoint, eight ML algorithms were developed to predict PASS status at the 24-month postoperative timepoint. The performance of each of these algorithms was evaluated, and its generalizability was assessed using a prospective internal test set.

Results The long short-term memory (LSTM)-based algorithm demonstrated the best performance (area under the receiver operating characteristic curve, 0.90 ± 0.13).

Conclusions The LSTM-based algorithm accurately predicted which group was likely to achieve PASS at the 24-month postoperative timepoint. Although this study included a small number of patients with limited available clinical data, the concept of using past outcomes to predict further outcomes presented herein may provide insights for optimizing clinical schedules and efficient medical resource utilization.

Trial registration This study was registered as a clinical trial (Clinical Trial No. NCT02487901), and the study protocol was approved by the Seoul National University Hospital Institutional Review Board (IRB No. 1505-037-670).

[†]Yechan Seo and Seoi Jeong contributed as first-authors.

*Correspondence:
Hyoun-Joong Kong
gongcop7@snu.ac.kr
Chi Heon Kim
chiheon1@snu.ac.kr

Full list of author information is available at the end of the article



© The Author(s) 2024. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

Keywords Cervical vertebra, Laminoplasty, Machine learning, Outpatient clinic, Patient outcome assessment, Telemedicine

Introduction

At predetermined intervals following spinal surgery, patients typically visit outpatient clinics for postoperative follow-up. Despite some variations, depending on the institution, doctor, and patient-specific condition, the postoperative follow-up typically includes visits at 1, 3, 6, and 12 months, with annual visits thereafter [1–9]. However, based on their perception that a routine visit is unnecessary, patients sometimes skip scheduled appointments. Moreover, the schedule is not usually individualized by clinical outcomes and is relatively inflexible. For example, despite symptomatic improvement, a patient may continue to adhere to a predetermined hospital visit schedule [1–12]. Systematic modulation of the outpatient visits to individualized clinical outcomes could improve the efficient use of medical resources, and thereby benefit both patients and healthcare providers through optimization of medical resource utilization. Previous algorithms for improving the efficiency of outpatient clinics prioritized patient status [13–19]. Research on clinical history-based prediction of future patient conditions is ongoing [20–23]. Predictive algorithms differ, based on the analytical method, such as machine learning (ML), which have been developed to predict postoperative states [24–30]. As these algorithms may not be applicable to every disease, the creation of customized algorithms for each condition is essential. For example, to predict the treatment outcomes of patients with lumbar disc herniation, Pedersen et al. recently used ML and deep learning by applying decision tree (DT), support vector machine (SVM), random forest (RF), and boosted tree algorithms [29].

Cervical laminoplasty, commonly used to treat cervical myelopathy [2, 4–9, 31–33], involves changes to the configuration of the spinal lamina to expand the cervical spinal canal, which is subsequently maintained using a metal plate/screw system and solid bony fusion at the lamina for approximately 1 year. Computed tomography is routinely performed at either 6 or 12 months postoperatively, and the stability of the internal structure is determined from the solid bony union that occurs at the hinge site [2, 8, 9]; thereafter, laminoplasty re-closure and the consequent neurological deterioration are unlikely. We hypothesized that a 2-year follow-up might not be essential for patients with symptomatic improvement and solid bony fusion. Nonetheless, as neurological symptoms may change for various reasons, we recommend the implementation of an alternative follow-up system, such as telemedicine.

To date, no study has successfully developed an ML model, based on long-term follow-up data (>2 years), for

the specific optimization of post-cervical laminoplasty outpatient scheduling in patients who had ossification of the posterior longitudinal ligament (OPLL)- or degenerative spinal disease-based cervical stenosis. To address this knowledge gap, in the present study, we enrolled a prospective cohort of patients who underwent cervical laminoplasty (Clinical Trial No. NCT02487901) and were scheduled to undergo follow-up at 1, 3, 6, 12, and 24 months postoperatively [2]. Using data from this prospective cohort, we conducted a pilot study to evaluate the feasibility of using the 12-month postoperative clinical outcomes to predict the 24-month postoperative outcomes. This study was conducted with an aim to develop clinical information-based predictive ML algorithms to stratify patients who would have stable outcomes at the 24-month postoperative timepoint and to identify the most appropriate algorithm for this purpose.

Methods

Study design and patient population

This post hoc analysis comprised a subgroup analysis of data from a prospective cohort study of 255 patients who, between July 2015 and April 2017, underwent cervical laminoplasty for OPLL- or degenerative spinal disease-induced cervical stenosis [2]. For all patients, the Arch™ laminoplasty system (DePuy Synthes, Oberdorf, Switzerland), with a 12-mm spacer length, was applied during cervical laminoplasty. All surgeries were performed by an experienced surgeon, with experience in conducting more than 500 cervical laminoplasty procedures over a decade. All surgeries were conducted with strict adherence to the standard cervical open laminoplasty procedures. All patients in the cohort were scheduled to visit the clinic at 1, 3, 6, 12, and 24 months postoperatively, at which point clinical outcomes, including the Japanese Orthopedic Association (JOA) scores, were prospectively collected [34]. This secondary analysis comprised data from 80 patients (M: F=48:32; age, 59.8 ± 10.1 years) who completed a 24-month follow-up schedule. To further validate the robustness and generalizability of our findings, we included a prospective internal test set with 22 additional patients, who underwent data collection between September 2020 and July 2022 and, to ensure consistency in patient selection and data collection, were recruited using the same methods as those used in the original cohort.

In this secondary analysis, data from a prospective cohort with cervical myelopathy were used to predict the patient acceptable symptom state (PASS) status at the 24-month postoperative timepoint. Therefore, we

excluded patients without the 24-month postoperative JOA data. Consequently, the final study sample comprised 80 of the 255 patients originally enrolled in the cohort. A flow diagram illustrating this process is shown in Fig. 1.

This study conformed to the principles evinced in the Declaration of Helsinki and the Guidelines for Good Clinical Practice. The Ethics Committee and the Institutional Review Board of Seoul National University Hospital approved (approval no. 1505-037-670) the study protocol and, owing to the retrospective nature of this study, waived the requirement of informed consent.

Variables and ML model implementation

The analysis included 17 clinikoradiological data points as independent variables, including age, sex, body mass index (BMI, kg/m²), diabetes, smoking status, occupation, Charlson Comorbidity Index (CCI), preoperative ambulatory status, diagnosis (stenosis vs. myelopathy), presence of high signal intensity on T2-weighted magnetic resonance imaging (HIS), presence of the snake eye sign (SES), ambulatory status at the 1-month postoperative timepoint, preoperative JOA score, and JOA scores at the 1-, 3-, 6-, and 12-month postoperative timepoints (Table 1). The JOA score assesses a patient's condition in six domains, on a scale from 0 to 17, with higher scores indicating a better patient condition [34]. The PASS was defined as a JOA score ≥ 14.25 at the 24-month postoperative timepoint [35]. Patients who met or did not meet the PASS criteria were coded as 1 and 0, respectively. A

PASS score of 1 indicated that the patient was in good condition. These coded values were used as dependent variables, wherein 63 (78.8%) and 17 (21.3%) patients were assigned codes of 1 and 0, respectively. Using the stratified random sampling technique, the training and test sets (8:2) were created; the training set comprised 50 and 14 patients with PASS 1 and 0 and the test set comprised 13 and 3 patients with PASS 1 and 0, respectively. Eight ML algorithms [36] were applied to the variables and dataset, as follows: logistic regression (LR) [37, 38], SVM [39], k-nearest neighbor (kNN) [40], RF [41], extreme gradient boosting (XGBoost) [42], multi-layer perceptron (MLP) [43, 44], recurrent neural network (RNN) [45], and long short-term memory (LSTM) [46]. LR is a representative ML algorithm that is broadly applied in various fields. This algorithm predicts a probability between 0 and 1 for PASS, the dependent variable, and is primarily used in binary classification. SVM is an algorithm that maximizes the distance (called a "margin") between two classes by using a linear kernel function to extend a linear space into a nonlinear space. kNN is one of the simplest available ML algorithms; with new input data, kNN identifies 'k' data points close to the existing set and subsequently classifies the data points as the class with the highest frequency of occurrence. We set three as the number of neighbors to consider (*k*). RF is an ensemble technique that combines multiple DTs to improve performance by using a bootstrap aggregation technique that generates a weak classifier for each sample by randomly extracting samples of the same size from




Fig. 1 The patients' flow diagram. The figure shows the overall flow of prospective study. Patients without JOA values at 24 months after surgery were excluded, and all other patients were included. Thus, 255 patients were enrolled in the prospective study, but 175 patients were excluded
Abbreviations: JOA: Japanese Orthopedic Association

Table 1 The demographics of patients

	Total (n = 80)	PASS = 1 (n = 63)	PASS = 0 (n = 17)	p value
Age (mean, std)	58.76 ± 10.10	57.05 ± 9.38	65.12 ± 10.41	0.008
Sex	Male :48 (60.0%) Female: 32 (40.0%)	Male: 36 (57.1%) Female: 27 (42.9%)	Male: 12 (70.6%) Female: 5 (29.4%)	0.315
BMI (mean, std)	25.78 ± 3.53	26.17 ± 3.42	24.32 ± 3.65	0.072
DM	Yes: 16 (20.0%) No: 64 (80.0%)	Yes: 13 (20.6%) No: 50 (79.4%)	Yes: 3 (17.6%) No: 14 (82.4%)	0.785
Smoking	Yes: 12 (15.0%) No: 68 (85.0%)	Yes :8 (12.7%) No: 55 (87.3%)	Yes: 4 (23.5%) No: 13 (76.5%)	0.267
OA ¹	1:6 (7.5%) 2:19 (23.8%) 3:55 (68.8%)	1:5 (7.9%) 2:15 (23.8%) 3:43 (68.3%)	1:1 (5.9%) 2:4 (23.5%) 3:12 (70.6%)	0.958
CCI (mean, std)	1.80 ± 1.30	1.62 ± 1.18	2.47 ± 1.50	0.042
Preop. Ambulation ²	1:19 (23.8%) 2:55 (68.8%) 3:6 (7.5%)	1:16 (25.4%) 2:41 (65.1%) 3:6 (9.5%)	1:3(17.6%) 2:14 (82.4%) 3:0 (0.0%)	0.281
Diagnosis	Stenosis: 3 (3.8%) Myelopathy: 77 (96.3%)	Stenosis: 1 (1.6%) Myelopathy: 62 (98.4%)	Stenosis: 2 (11.8%) Myelopathy: 15 (88.2%)	0.050
HIS	Yes: 77 (96.3%) No: 3 (3.8%)	Yes: 60 (95.2%) No: 3 (4.8%)	Yes: 17 (100.0%) No: 0 (0.0%)	0.359
SES	Yes: 24 (30.0%) No: 56 (70.0%)	Yes: 19 (30.2%) No: 44 (69.8%)	Yes: 5 (29.4%) No: 12 (70.6%)	0.952
Postop. 1 month ambulation ³	Nulls: 3 (3.8%) 1:28 (35.0%) 2:46 (57.5%) 3:3 (3.8%)	Nulls: 3 (4.8%) 1:23 (36.5%) 2:34 (54.0%) 3:3 (4.8%)	Nulls: 0 (0.0%) 1:5 (29.4%) 2:12 (70.6%) 3:0 (0.0%)	0.484
JOA score preop (mean, std)	11.18 ± 3.28	11.90 ± 2.77	8.47 ± 3.66	0.002
1 month (mean, std)	14.10 ± 2.21	14.60 ± 1.73	12.24 ± 2.80	0.003
3 months (mean, std)	14.92 ± 1.76	15.37 ± 1.38	13.24 ± 2.02	0.001
6 months (mean, std)	15.01 ± 1.75	15.56 ± 1.34	13.00 ± 1.66	0.000
12 months (mean, std)	15.23 ± 1.75	15.83 ± 0.99	13.00 ± 2.15	0.000
24 months (mean, std)	15.14 ± 2.42	16.08 ± 0.81	12.65 ± 1.50	0.000

¹ Occupational activity; 1: high, 2: intermediate, 3: low

² Preop. Ambulation; 1: fully ambulant, 2: ambulant with an aid, 3: no outdoor self-walking

³ Postop. 1 month ambulation; Null: no data, 1: fully ambulant, 2: ambulant with an aid, 3: no outdoor self-walking

Abbreviations: BMI: body mass index; CCI: Charlson comorbidity index; DM: diabetes mellitus; HIS: presence of high signal intensity in T2-weighted magnetic resonance imaging; OA: Occupational Activity; PASS: patient acceptable symptom state; SES: presence of snake eye sign; std: standard deviation

the original dataset multiple times. We used 100 trees, the two minimum numbers of samples required to split an internal node, one minimum node size, and the Gini Impurity Index as the node-splitting criteria. XGBoost is an algorithm of the ensemble-boosting technique that inputs errors between the actual and predicted values from previous models as training data and supplements errors using gradients. We used 400 trees, 3 maximum depths, and a 0.1 learning rate. An MLP is a neural network wherein one or more perceptrons form multiple layers, with one or more hidden layers between the input

and output layers. The algorithm was developed using stochastic gradient descent (SGD) for weight optimization, and the strength of L2 regularization was set at 1e-5 alpha, with a learning rate of 0.001, 32 neurons in three hidden layers, and a rectified linear unit (ReLU) function. An RNN is defined as an artificial neural network that is characterized by a recurrent structure that processes the inputs and outputs in sequential units. For the RNN model, 32, 10, and 2 hidden state sizes were defined, and the activation function was softmax. Categorical cross-entropy and Adam were used as the loss function and

optimizer, respectively. LSTM is an algorithm that was originally developed as one of the RNN algorithms to prevent the vanishing-gradient problem of existing RNNs by using cells, input gates, outputs, and forget gates. The LSTM uses the same hyperparameter settings as the RNN mentioned above. The detailed hyperparameters for each of the eight ML algorithms are listed in Table 2.

Feature engineering

Patient characteristics, including age, sex, diagnosis, BMI, diabetes, smoking status, and comorbidities, were obtained from the patients' nursing records. The presence of HIS and SES on T2-weighted magnetic resonance images was assessed, and radiology medical records were referenced. Sex was coded as 1 for male and 0 for female patients, whereas age was recorded as an integer. The diagnosis was coded as 1 and 2, respectively, for stenosis without and with clinical myelopathy, which included an increased deep tendon reflex, positive Hoffmann's sign, decreased grip and release test, positive Romberg test, or veering on tandem gait. The presence of SES and HIS was coded as 1 if yes and 0 if no. BMI was recorded as a continuous variable. Diabetes was coded as 1 if present, and 0 otherwise. The smoking status was coded as 1 for yes and 0 for no. Occupational classification was

undertaken according to the Occupational Activity (OA) criteria established by Steeves et al. [47], with high, intermediate, and low OA coded as 1, 2, and 3, respectively. The CCI is a validated method that, based on the presence and severity of 17 specific comorbidities, assigns a weighted score to predict long-term mortality and morbidity [48]. The preoperative and 1-month postoperative ambulatory statuses were coded as 1, 2, and 3 for fully possible ambulation, walking with an aid (stick, walker, etc.), and not being able to walk unaided outside, respectively. Patient prognosis was measured using the JOA pain score (from 0 [worst] to 17 [best]). The dependent variable of PASS status was defined as 1 or 0 if the JOA value at the 24-month postoperative timepoint was ≥ 14.25 and < 14.25 , respectively. To eliminate missing values during the 12-month follow-up period, which were imputed using linear interpolation, pre-processing was performed. Missing values for the categorical variables were imputed using a one-hot encoding method. However, patients with missing JOA values the 24-month postoperative timepoint were excluded from the training data to avoid errors in the ground truth.

Table 2 Hyperparameters of ML algorithms. This table summarizes the key hyperparameters used for various machine learning algorithms in the study. The hyperparameters were tuned and optimized to improve the model performance

Algorithms	Hyperparameter	Value
LR	-	-
SVM	Kernel	Linear
kNN	Number of neighbors (k)	3
RF	Number of trees	100
	Minimum samples to split a node	2
	Minimum node size	1
	Node splitting criterion	Gini Impurity
XGBoost	Number of trees	400
	Maximum depth	3
	Learning rate	0.1
MLP	Optimization algorithm	Stochastic Gradient Descent (SGD)
	L2 regularization strength	1e-5
	Learning rate	0.001
	Number of hidden layers	3
	Number of neurons per hidden layer	32
	Activation function	Rectified Linear Unit (ReLU)
RNN	Hidden state sizes	30, 10, 2
	Activation function	Softmax
	Loss function	Categorical cross-entropy
	Optimizer	Adam
LSTM	Hidden state sizes	30, 10, 2
	Activation function	Softmax
	Loss function	Categorical cross-entropy
	Optimizer	Adam

Abbreviations: ML: machine learning; LR: Logistic regression; SVM: Support vector machines; kNN: k-nearest neighbor; XGBoost: Extreme gradient boosting; RF: Random forests; MLP: Multilayer perceptron; RNN: Recurrent neural network; LSTM: Long short-term memory

Statistical analysis

Table 3 presents a comparison of the excluded and included populations. To assess the potential selection bias, we analyzed intergroup differences in the 17 independent variables. For numerical variables, such as age, BMI, CCI, and JOA scores, we conducted independent sample *t*-tests, which were preceded by Levene's test for equality of variances to verify the assumption of intergroup homogeneity of variance. When Levene's test

indicated equal or unequal variances ($p > 0.05$ or $p \leq 0.05$, respectively), a standard *t*-test or Welch's *t*-test (to adjust for this difference), was performed. For categorical variables, including sex, DM, smoking status, OA, preoperative ambulation, postoperative ambulation, diagnosis, residence, HIS, and SES, a cross-tabulation analysis, followed by a chi-square test, was conducted to determine the intergroup distributional differences. To ensure consistent and robust comparisons between the PASS 1 and

Table 3 Differences in demographics between included and excluded patients

	Total (n = 255)	Included (n = 80)	Excluded (n = 175)	p value
Age (mean, std)	59.64 ± 12.14	58.76 ± 10.10	60.03 ± 12.97	0.396
Sex	Male: 181 (71.0%) Female: 74 (29.0%)	Male: 48 (60.0%) Female: 32 (40.0%)	Male: 133 (76.0%) Female: 42 (24.0%)	0.009
BMI (mean, std)	25.37 ± 3.71	25.78 ± 3.53	25.19 ± 3.78	0.226
DM	Yes: 55 (21.6%) No: 200 (78.4%)	Yes: 16 (20.0%) No: 64 (80.0%)	Yes: 39 (22.3%) No: 136 (77.7%)	0.681
Smoking	Yes: 50 (19.6%) No: 205 (80.4%)	Yes: 12 (15.0%) No: 68 (85.0%)	Yes: 38 (21.7%) No: 137 (78.3%)	0.210
OA ¹	1: 20 (7.8%) 2: 44 (17.3%) 3: 191 (74.9%)	1: 6 (7.5%) 2: 19 (23.8%) 3: 55 (68.8%)	1: 14 (8.0%) 2: 25 (14.3%) 3: 136 (77.7%)	0.178
CCI (mean, std)	1.92 ± 1.39	1.80 ± 1.30	1.97 ± 1.44	0.345
Preop. Ambulation ²	Nulls: 1 (0.4%) 1: 74 (29.0%) 2: 153 (60.0%) 3: 27 (10.6%)	1: 19 (23.8%) 2: 55 (68.8%) 3: 6 (7.5%)	Nulls: 1 (0.6%) 1: 55 (31.4%) 2: 98 (56.0%) 3: 21 (12.0%)	0.162
Diagnosis	Stenosis: 14 (5.5%) Myelopathy: 241 (94.5%)	Stenosis: 3 (3.8%) Myelopathy: 77 (96.3%)	Stenosis: 11 (6.3%) Myelopathy: 164 (93.7%)	0.409
HIS	Yes: 239 (93.7%) No: 16 (6.3%)	Yes: 77 (96.3%) No: 3 (3.8%)	Yes: 162 (92.6%) No: 13 (7.4%)	0.261
SES	Yes: 91 (35.7%) No: 164 (64.3%)	Yes: 24 (30.0%) No: 56 (70.0%)	Yes: 67 (38.3%) No: 108 (61.7%)	0.200
Postop. 1 month ambulation ³	Nulls: 10 (3.9%) 1: 102 (40.0%) 2: 129 (50.6%) 3: 14 (5.5%)	Nulls: 3 (3.8%) 1: 28 (35.0%) 2: 46 (57.5%) 3: 3 (3.8%)	Nulls: 7 (4.0%) 1: 74 (42.3%) 2: 83 (47.4%) 3: 11 (6.3%)	0.291
JOA score preop (mean, std)	11.10 ± 3.49	11.18 ± 3.28	11.07 ± 3.59	0.816
1 month (mean, std)	13.88 ± 2.60	14.10 ± 2.21	13.78 ± 2.76	0.321
3 months (mean, std)	14.72 ± 2.04	14.92 ± 1.76	14.61 ± 2.19	0.257
6 months (mean, std)	14.87 ± 2.06	15.01 ± 1.75	14.78 ± 2.24	0.408
12 months (mean, std)	15.16 ± 1.96	15.23 ± 1.75	15.10 ± 2.14	0.228
Residence ⁴	Nulls: 16 (6.3%) 1: 106 (41.6%) 2: 54 (21.2%) 3: 79 (31.0%)	Nulls: 6 (7.5%) 1: 31 (38.8%) 2: 24 (30.0%) 3: 19 (23.8%)	Nulls: 10 (5.7%) 1: 75 (42.9%) 2: 30 (17.1%) 3: 60 (34.3%)	0.080

¹ Occupational activity; 1: high, 2: intermediate, 3: low

² Preop. Ambulation; Null: no data, 1: fully ambulant, 2: ambulant with an aid, 3: no outdoor self-walking

³ Postop. 1 month ambulation; Null: no data, 1: fully ambulant, 2: ambulant with an aid, 3: no outdoor self-walking

⁴ Residence; Null: no data, 1: live in Seoul, 2: live in Seoul Metropolitan Area except for Seoul (Incheon, Gyeonggi), 3: live outside the Seoul Metropolitan Area

0 groups, the same statistical methods were applied to analyze the variables that have been presented in Table 1.

Evaluation of the ML model

The output of the analysis was the PASS score at the 24-month postoperative timepoint. Eight ML algorithms were applied to the data: LR, SVM, kNN, RF, XGBoost, MLP, RNN, and LSTM. The performance of each algorithm was evaluated from the sensitivity, specificity, positive predictive value (PPV), negative predictive value (NPV), F1-score, accuracy, and area under the receiver operating characteristic curve (AUROC) [49]. To address the issue of a small number of test samples, the k-fold cross-validation method was applied, and the 5-fold cross validation was ultimately used in this study. Additionally, the robustness and generalizability of the model were validated using a prospective internal test set, and the feature importance of the best-performing model was analyzed using Shapley additive explanation (SHAP) values [50].

This study was conducted using Python version 3.6.13, XGBoost version 1.6.1, SciPy version 1.8.1, Scikit-learn version 1.1.1, Seaborn version 0.11.2, Pandas version 1.4.3, NumPy version 1.19.5, TensorFlow version 2.0.0, and Keras version 2.3.1. Statistical analysis was carried out from July 26, 2022, to August 2, 2022.

Results

Patient outcomes

The patient demographics are presented in Table 1. Pre-operatively, the mean JOA score was 11.18 ± 3.28 , which significantly improved to 14.10 ± 2.21 at 1 through 24 months postoperatively ($p < 0.05$, Table 1) and was maintained for 24 months. A significant intergroup difference was observed between PASS 1 and 2 groups at each timepoint ($p < 0.05$, as detailed in Supplementary Material 1).

Algorithm performance

The performances of the eight ML algorithms are presented in Table 4. For each evaluation metric, the performance was analyzed five times using 5-fold cross

validation. In terms of the average values, the algorithm with the best performance for each metric, based on the specificity, sensitivity, PPV, NPV, F1-score, accuracy, and AUROC, was LSTM (specificity, 0.883 ± 0.211 ; sensitivity, 0.967 ± 0.041 ; PPV, 0.955 ± 0.061 ; NPV, 0.893 ± 0.137 ; F1-score, 0.960 ± 0.044 ; accuracy, 0.938 ± 0.069 ; and AUROC, 0.900 ± 0.130114). A temporal external test of the LSTM-based algorithm was conducted on the test set comprising 22 patients. This model was selected because of its excellent performance and generated the following results from the prospective internal test set: specificity, 0.892 ± 0.062 ; sensitivity, 0.875 ± 0.0 ; PPV, 0.84 ± 0.07 ; NPV, 0.92 ± 0.006 ; F1-score, 0.856 ± 0.039 ; accuracy, 0.886 ± 0.038 ; and AUROC, 0.858 ± 0.007 . Figure 2 shows the receiver operating characteristic (ROC) curves for the LSTM-based algorithm, where (a) represents the ROC curve for the internal test set and (b) represents the ROC curve for the prospective internal test set. Figure 3 shows the ROC curves of the eight ML models. The threshold for all ROC curves was set to 0.5. This figure provides a visual comparison of the performance of each model for distinguishing between patients with stable outcomes and those who require further follow-up. For more detailed information, refer to Figs. S1–S7. Figure 4 shows the feature importance of all 17 independent variables in the top-performing LSTM-based algorithm, as determined from the SHAP values.

Discussion

This study aimed to evaluate the predictability of the 2-year postoperative outcomes based on the 1-year postoperative outcomes. To achieve this, various ML algorithms were tested, with the LSTM exhibiting the best average performance. The high specificity (0.883 ± 0.211) of this LSTM-based algorithm indicates that it could accurately identify patients who would need to visit the outpatient clinic (i.e., patients with PASS=0) at the 24-month postoperative timepoint. Despite using a relatively small training set, the algorithm demonstrated strong performance on a prospective internal test set, and this supports its generalizability. These results may

Table 4 The performance of the eight ML algorithms

ML Model	Sensitivity	Specificity	PPV	NPV	F1-Score	Accuracy	AUROC
LR	0.91 ± 0.03	0.52 ± 0.17	0.88 ± 0.03	0.58 ± 0.15	0.89 ± 0.03	0.83 ± 0.05	0.71 ± 0.09
SVM	0.78 ± 0.15	0.65 ± 0.23	0.90 ± 0.06	0.49 ± 0.11	0.82 ± 0.08	0.75 ± 0.08	0.71 ± 0.05
kNN	0.94 ± 0.06	0.47 ± 0.25	0.87 ± 0.05	0.67 ± 0.37	0.90 ± 0.04	0.84 ± 0.08	0.70 ± 0.14
RF	0.92 ± 0.12	0.58 ± 0.33	0.90 ± 0.06	0.62 ± 0.38	0.90 ± 0.05	0.85 ± 0.06	0.75 ± 0.13
XGBoost	0.94 ± 0.06	0.70 ± 0.40	0.93 ± 0.08	0.63 ± 0.34	0.93 ± 0.02	0.89 ± 0.05	0.82 ± 0.18
MLP	0.95 ± 0.06	0.53 ± 0.20	0.88 ± 0.05	0.80 ± 0.27	0.92 ± 0.04	0.86 ± 0.07	0.74 ± 0.12
RNN	0.95 ± 0.04	0.77 ± 0.29	0.94 ± 0.07	0.79 ± 0.19	0.95 ± 0.05	0.91 ± 0.09	0.86 ± 0.16
LSTM	0.97 ± 0.04	0.88 ± 0.21	0.96 ± 0.06	0.89 ± 0.14	0.96 ± 0.04	0.94 ± 0.07	0.90 ± 0.13

Abbreviations: AUROC: area under the receiver operating characteristic; ML: machine learning; NPV: negative predictive value; PPV: positive predictive value; LR: Logistic regression; SVM: Support vector machines; kNN: k-nearest neighbor; XGBoost: Extreme gradient boosting; RF: Random forests; MLP: Multilayer perceptron; RNN: Recurrent neural network; LSTM: Long short-term memory




Fig. 2 ROC curves of the LSTM-based algorithm. **(a)** ROC curve for the internal test set. **(b)** ROC curve for the prospective internal test set. Due to the 5-fold cross-validation, five ROC curves are generated for each test set, with the mean ROC curve (blue line) representing the average performance of the five models. The shaded area indicates the standard deviation across the folds. Although the AUROC value is lower for the prospective internal test set (0.86 ± 0.01) compared to the internal test set (0.90 ± 0.13), the LSTM-based model still demonstrates strong performance

Abbreviations: ROC: Receiver-operating characteristic; AUROC: Area under the receiver operating characteristic; LSTM: Long short-term memory

be useful for modifying the post-cervical laminoplasty clinical follow-up schedule.

Inter-model comparison

In this study, we tested five classical ML algorithms (LR, SVM, kNN, RF, and XGBoost) as well as three deep-learning algorithms (MLP, RNN, and LSTM), of which the RNN- and LSTM-based algorithms demonstrated a higher average performance across all metrics than the classical ML algorithms. The superior performance of time series-based deep-learning models, such as LSTM and RNN, could be attributed to their ability to effectively process sequential data [51, 52]. In this context, the JOA values were ordered chronologically, indicating that past patient outcomes significantly influenced future outcomes, and thereby makes these models particularly well-suited for this task.

As shown in Fig. 3, the ROC curve for the LSTM-based algorithm was skewed more toward the top-left corner than in any of the other models, which resulted in an AUROC value > 0.9 that indicates excellent performance. In the present study, the high specificity (0.883 ± 0.211) of the model was considered particularly important as it accurately identifies patients who need to visit an outpatient clinic at 2 years postoperatively (i.e., those with PASS 0). Therefore, we concluded that the LSTM-based model was the most suitable for this purpose.

Feature importance

By calculating the impact of each feature on the predicted outcome, the SHAP values for the LSTM-based algorithm (Fig. 4) provide insights into how each feature contributes to the model's predictions. The top 10 features, ranked by SHAP values, include the JOA scores at the 12-, 6-, 1-, and 3-month postoperative timepoints; BMI; preoperative JOA score; sex; age; SES; and preoperative ambulatory status. Notably, BMI exhibited a negative correlation with predicted outcomes, whereas sex showed a negative correlation for male patients and a positive correlation for female patients. Specifically, the JOA score at the 1-month postoperative timepoint negatively correlated, whereas the JOA scores at the 3-, 6-, and 12-month postoperative timepoints positively correlated, with the model predictions. This underscores the significance of time-series data up to the 12-month postoperative timepoint for predicting the 2-year postoperative outcomes. Detailed SHAP values for each fold are shown in Figs. S8–S12.

The high SHAP values of the JOA scores indicate that these sequential clinical outcomes are crucial for determining whether a patient will achieve satisfactory results at the 24-month postoperative timepoint. This result aligns with the ability of the LSTM-based algorithm to effectively capture the temporal progression of patient outcomes. Furthermore, clinical information such as age




Fig. 3 Mean ROC curves for the eight ML models. The plot shows the mean ROC curves for eight ML algorithms: LR, SVM, kNN, RF, XGBoost, MLP, RNN, and LSTM. The AUROC values for each model, along with their standard deviations, are displayed in the legend. The LSTM-based algorithm demonstrates the highest mean AUROC (0.9 ± 0.13), indicating superior performance compared to the other models

Abbreviations: ROC: Receiver-operating characteristic; AUROC: Area under the receiver operating characteristic; LR: Logistic regression; SVM: Support vector machines; kNN: k-nearest neighbor; XGBoost: Extreme gradient boosting; RF: Random forests; MLP: Multilayer perceptron; RNN: Recurrent neural network; LSTM: Long short-term memory; ML: Machine learning

and SES, contributed significantly, and this underscores the model's capacity to integrate both time-dependent and static variables for accurate prediction.

This detailed analysis confirmed the clinical relevance of the features selected for the LSTM-based algorithm, and thereby enable more informed clinical decision-making by using both time-series data and clinical information.

Alternative approaches in resource-limited settings

If a hospital environment lacks the computational resources required to deploy an LSTM-based algorithm for inference, a classical ML model may be a suitable alternative. In this case, the XGBoost model is recommended as the next-best option because it demonstrates the highest specificity among the five classical ML models.

Additionally, it may be challenging to collect all the 17 clinicoradiological values introduced in this study for all outpatients in a hospital setting. In such cases, variables with the highest SHAP values (e.g., JOA scores up to 12 months postoperatively), as shown in Fig. 4, should be prioritized for data collection.

Clinical application

In patients who underwent cervical laminoplasty, the patient's condition stabilized following solid bony fusion at the hinge of the reflected lamina, which occurred between 6 and 12 months [2, 8, 9, 32]. Therefore, if a patient's condition stabilizes for a year, it is unlikely to worsen subsequently [2, 8, 9]. These results indicate that patients who achieve stable clinical improvement may not need to visit outpatient clinics frequently. Given these findings, it would be practical to develop an



Fig. 4 Shapley additive explanation (SHAP) values of all 17 independent variables in the top-performing LSTM-based algorithm
Abbreviations: SHAP: Shapley additive explanation; LSTM: Long short-term memory

LSTM-based algorithm that uses the clinicoradiological data from the first postoperative year to identify patients who may safely forego the 2-year postoperative follow-up clinic visit. The identification of patients who meet the PASS criteria could reduce the need for routine in-person follow-up visits and thereby optimize the use of limited medical resources. The algorithm developed in this study could be integrated into an electronic medical information system to assist decision-making for clinical follow-up scheduling. However, this algorithm is not perfect, and this may have led to patients missing clinic visits despite worsening clinical outcomes [1–8, 10–12, 32]. Therefore, this algorithm should not be used to completely exempt patients from clinical visits. Instead, it can

be employed to modify the visit frequency or guide the incorporation of alternative follow-up methods, such as telemedicine [53–55]. However, it is crucial to note that, although telemedicine offers a viable option for reducing the number of in-person visits, particularly for patients with a PASS 1, to ensure that patient safety is prioritized, it should be considered an adjunct, rather than a replacement, of direct clinical assessment.

Research significance and novelty

Our study builds on prior research that highlights the importance of predicting patient no-shows and optimizing outpatient schedules, based on patient characteristics, and minimizing unnecessary clinic visits to

efficiently use medical resources [13, 15–19, 56]. Importantly, this novel study applied an ML-based approach to optimize outpatient follow-up schedules, specifically for patients with cervical myelopathy after cervical laminoplasty, with follow-up >2 years. Although ML models have been previously applied to optimize outpatient scheduling for other patient populations, such as those with lumbar disc herniation [29], no study had specifically developed post-laminoplasty schedules for patients with cervical myelopathy by using long-term (>2-year) follow-up data. This highlights the uniqueness and significance of our study in addressing this clinical need.

In addition, similar to prior research that demonstrated temporal generalization using prospective internal test sets in different periods [57–59], our LSTM-based algorithm successfully underwent temporal generalization validation using a prospective internal test set. As shown in Fig. 2, although the ROC curve for the prospective internal test set exhibited a slightly lower AUROC value compared to the internal test set, its specificity (0.892 ± 0.062) was higher than that of the internal test set (0.883 ± 0.211) and indicated that our model is not overfitted to a specific cohort, but rather, demonstrates generalizability across different periods.

Variables utilized for ML

The current study used the JOA score as the most important clinical outcome measure in patients with cervical myelopathy. Although the JOA is widely used to measure the post-cervical laminoplasty clinical outcome, it may not sufficiently represent a patient's status. The inclusion of additional variables, such as quality of life and numeric rating pain scores for the neck and arm, would therefore help to improve the performance of the algorithm. Defining the PASS by using multiple variables may further improve the reliability of the ML algorithm; however, this proposal was not explored in the present study, and should be further investigated in future studies.

Limitations

This study had some limitations. First, a small number of data points were used for the training set. Besides the limited sample size, the study population was drawn exclusively from a single institution – Seoul National University Hospital – that primarily serves Korean patients, and this resulted in a lack of ethnic diversity. The data collection period was limited, which may have restricted the generalizability of the findings. Although clinical information was collected prospectively from 255 patients, owing to missing datapoints, data from only 80 patients were usable. A five-fold cross-validation algorithm was applied to address the issue of the small sample. Furthermore, to demonstrate its generalizability, the algorithm was validated using a prospective internal

test set that was collected at a significantly different period, which highlights its robustness across temporal variations. However, before general clinical use, the algorithm should be tailored and optimized by analyzing large datasets. Further research using large and complete datasets from various institutions, and including patients of diverse ethnic background, is necessary to optimize the algorithm. A prospective cohort study to evaluate the clinical utility and generalizability of this model is required. Second, the PASS criteria of the JOA may not fully represent the patient's status. For example, depending only on the clinical outcomes, OPLL progression may be missed. Further research is required to address this issue and to accurately identify patients who may not require in-person clinic visits. Despite these limitations, this study is the first to apply ML algorithms to predict the post-cervical laminoplasty 24-month postoperative outcomes using data from a prospective cohort. This information could enable the development of algorithms to modify the clinic-visit schedule or the type of clinic.

Conclusions

Despite the small sample size, using clinicoradiological data from a prospective cohort, this study demonstrated the robust predictive performance of an ML algorithm for the 24-month postoperative outcome. Based on their clinical data for up to 1 year, the predictive identification of patients who would achieve stable outcomes at the 24-month postoperative timepoint could facilitate more efficient medical resource utilization.

Abbreviations

AUROC	Area under the receiver operating characteristic
BMI	Body mass index
CCI	Charlson Comorbidity Index
DM	Diabetes mellitus
HIS	Presence of high signal intensity in T2-weighted magnetic resonance imaging
ML	Machine learning
NPV	Negative predictive value
OA	Occupational Activity
PASS	Patient acceptable symptom state
PPV	Positive predictive value
SES	Presence of snake eye sign
Std	Standard deviation

Supplementary Information

The online version contains supplementary material available at <https://doi.org/10.1186/s12911-024-02693-y>.

Supplementary Material 1

Supplementary Material 2

Acknowledgements

This work was supported by the New Faculty Startup Fund from Seoul National University. This study was supported by grant no. 3020230120 from Seoul National University Hospital research fund. This study was supported by Doosan Yonkang foundation (800-20210527). This research was

supported by a grant (2023MDD0075) from Armed Forces Capital Hospital in 2023 (800-20230466). This research was supported by a grant of 'Korea Government Grant Program for Education and Research in Medical AI' through the Korea Health Industry Development Institute (KHIDI), funded by the Korea government (MOE, MOHW). This study was supported by a grant from the National R&D Program for Cancer Control, Ministry of Health & Welfare, Republic of Korea (RS-2023-CC140357).

Author contributions

Yechan Seo: YS, Seoi Jeong: SJ, Sungwan Kim: SK, Siyoung Lee: SL, Jun-Hoe Kim: JHK, Tae-Shin Kim: TSK, Chun Kee Chung: CKC, Chang-Hyun Lee: CHL, John M. Rhee: JMR, Hyoun-Joong Kong: HJK, Chi Heon Kim: CHK. Conception and design: YS, SJ, HJK, CHK. Acquisition and data: YS, SJ, JHK, TSK, CKC, CHK, CHL. Analysis and interpretation of data: YS, SJ, SK, JMR, HJK, CHK. Drafting of the manuscript: YS, SJ, SK, JMR, HJK, CHK, CKC, SL. Critical revision of the manuscript: SK, CHK, HJK, JMR. Statistical analysis: YS, SJ, SK, HJK. Obtaining funding: CHK. Administrative, technical and material support: HJK, CHK. Supervision: SK, JMR, CKC.

Funding

This work was supported by the New Faculty Startup Fund from Seoul National University. This study was supported by grant no. 3020230120 from Seoul National University Hospital research fund. This study was supported by Doosan Yonkang foundation (800-20210527). This research was supported by a grant (2023MDD0075) from Armed Forces Capital Hospital in 2023 (800-20230466). This research was supported by a grant of 'Korea Government Grant Program for Education and Research in Medical AI' through the Korea Health Industry Development Institute (KHIDI), funded by the Korea government (MOE, MOHW). This study was supported by a grant from the National R&D Program for Cancer Control, Ministry of Health & Welfare, Republic of Korea (RS-2023-CC140357).

Data availability

The materials and data used in this study will be shared upon reasonable written request to the corresponding author. Access to the code requires the data access agreement and permission from the institutional review board.

Declarations

Ethics approval and consent to participate

This study was conducted in accordance with the Declaration of Helsinki and the Guideline for Good Clinical Practice. The study protocol was approved by the Seoul National University Hospital ethics committee/institutional review board (1505-037-670). The Seoul National University Hospital ethics committee/institutional review board approved the exemption of informed consent due to the retrospective nature of this study.

Consent for publication

Not applicable.

Competing interests

The authors declare no competing interests.

Author details

¹Department of Medicine, Seoul National University College of Medicine, 103 Daehak-ro, Jongno-gu, Seoul 03080, Republic of Korea

²Department of Transdisciplinary Medicine, Seoul National University Hospital, 101 Daehak-Ro, Jongno-gu, Seoul 03080, Republic of Korea

³Innovative Medical Technology Research, Seoul National University Hospital, 101 Daehak-Ro, Jongno-gu, Seoul 03080, Republic of Korea

⁴School of Medicine, the Faculty of Medical Science, Newcastle University, Newcastle Upon Tyne NE2 4HH, UK

⁵Department of Neurosurgery, Champodonamu Hospital, 32 Baumoe-ro 35-gil, Seocho-gu, Seoul 03080, Republic of Korea

⁶Department of Neurosurgery, Seoul National University Hospital, 101, Daehak-ro, Jongno-gu, Seoul 03080, Republic of Korea

⁷Department of Neurosurgery, Seoul National University College of Medicine, 103 Daehak-ro, Jongno-gu, Seoul 03080, Republic of Korea

⁸Department of Brain and Cognitive Sciences, Seoul National University, 101, 1, Gwanak-ro, Gwanak-gu, Seoul, Republic of Korea

⁹Department of Orthopaedic Surgery, Emory University School of Medicine, Atlanta, GA 30322, USA

¹⁰Department of Medical Device Development, Seoul National University College of Medicine, 103 Daehak-ro, Jongno-gu, Seoul 03080, Republic of Korea

Received: 22 January 2024 / Accepted: 24 September 2024

Published online: 30 September 2024

References

1. Rigal J, Quarto E, Boue L, Balabaud L, Thompson W, Cloche T, et al. Original surgical treatment and long-term follow-up for chronic inflammatory demyelinating polyradiculoneuropathy causing a compressive cervical myelopathy: review of the literature. *Neurospine*. 2022;19(2):472–7.
2. Kim CH, Chung CK, Choi Y, Kuo CC, Lee U, Yang SH, et al. The efficacy of ultrasonic bone scalpel for unilateral cervical open-door laminoplasty: a randomized controlled trial. *Neurosurgery*. 2020;86(6):825–34.
3. Riew KD. Double dome laminoplasty: works well but there are exceptions. *Neurospine*. 2021;18(4):889–90.
4. Ono K, Murata S, Matsushita M, Murakami H. Cervical lordosis ratio as a novel predictor for the loss of cervical lordosis after laminoplasty. *Neurospine*. 2021;18(2):311–8.
5. Nagoshi N, Nori S, Tsuji O, Suzuki S, Okada E, Yagi M, et al. Surgical and functional outcomes of expansive open-door laminoplasty for patients with mild kyphotic cervical alignment. *Neurospine*. 2021;18(4):749–57.
6. Lee DH, Dadufalza GKP, Baik JM, Park S, Cho JH, Hwang CJ, Lee CS. Double dome laminoplasty: a novel technique for C2 decompression. *Neurospine*. 2021;18(4):882–8.
7. Brown NJ, Lien BV, Shahrestani S, Choi EH, Tran K, Gattas S, et al. Getting down to the bare bones: does laminoplasty or laminectomy with fusion provide better outcomes for patients with multilevel cervical spondylotic myelopathy? *Neurospine*. 2021;18(1):45–54.
8. Lee S, Chung CK, Kim CH. Risk factor analysis of hinge fusion failure after plate-only open-door laminoplasty. *Global Spine J*. 2015;5(1):9–16.
9. Lee SE, Chung CK, Kim CH, Jahng TA. Symmetrically medial bony gutters for open-door laminoplasty. *J Spinal Disord Tech*. 2013;26(3):E101–6.
10. Kim N, Kim TH, Oh JK, Lim J, Lee KU, Kim SW. Analysis of the incidence and risk factors of postoperative delirium in patients with degenerative cervical myelopathy. *Neurospine*. 2022;19(2):323–33.
11. Kim JY, Hong HJ, Lee DC, Kim TH, Hwang JS, Park CK. Comparative analysis of 3 types of minimally invasive posterior cervical foraminotomy for foraminal stenosis, uniportal-, biportal endoscopy, and microsurgery: radiologic and midterm clinical outcomes. *Neurospine*. 2022;19(1):212–23.
12. Kim JY, Heo DH, Lee DC, Kim TH, Park CK. Comparative analysis with modified inclined technique for posterior endoscopic cervical foraminotomy in treating cervical osseous foraminal stenosis: radiological and midterm clinical outcomes. *Neurospine*. 2022;19(3):603–15.
13. Creps J, Lotfi V. A dynamic approach for outpatient scheduling. *J Med Econ*. 2017;20(8):786–98.
14. Hendrickson SB, Sims NM, DaSilva KA, Vallier HA. Improvement in outpatient follow-up with a postdischarge phone call intervention. *J Am Acad Orthop Surg*. 2020;28(18):e815–22.
15. Idowu OA, Boyajian HH, Ramos E, Shi LL, Lee MJ. Trend of spine surgeries in the outpatient hospital setting versus ambulatory surgical center. *Spine (Phila Pa 1976)*. 2017;42(24):E1429–36.
16. Kyriacou DN, Handel D, Stein AC, Nelson RR. BRIEF REPORT: factors affecting outpatient follow-up compliance of emergency department patients. *J Gen Intern Med*. 2005;20(10):938–42.
17. Ogulata SN, Cetik MO, Koyuncu E, Koyuncu M. A simulation approach for scheduling patients in the department of radiation oncology. *J Med Syst*. 2009;33(3):233–9.
18. Philpott-Morgan S, Thakrar DB, Symons J, Ray D, Ashrafian H, Darzi A. Characterising the nationwide burden and predictors of unkept outpatient appointments in the National Health Service in England: a cohort study using a machine learning approach. *PLoS Med*. 2021;18(10):e1003783.
19. Vermeulen IB, Bohte SM, Elkhuisen SG, Lameris H, Bakker PJ, La Poutre H. Adaptive resource allocation for efficient patient scheduling. *Artif Intell Med*. 2009;46(1):67–80.

20. Costelloe C, Burns S, Yong RJ, Kaye AD, Urman RD. An analysis of predictors of persistent postoperative pain in spine surgery. *Curr Pain Headache Rep*. 2020;24(4):11.
21. Manz CR, Chen J, Liu M, Chivers C, Regli SH, Braun J, et al. Validation of a machine learning algorithm to predict 180-day mortality for outpatients with cancer. *JAMA Oncol*. 2020;6(11):1723–30.
22. Schröder ML, de Wispelaere MP, Staartjes VE. Predictors of loss of follow-up in a prospective registry: which patients drop out 12 months after lumbar spine surgery? *Spine J*. 2019;19(10):1672–9.
23. Veeravagu A, Li A, Swinney C, Tian L, Moraff A, Azad TD, et al. Predicting complication risk in spine surgery: a prospective analysis of a novel risk assessment tool. *J Neurosurg Spine*. 2017;27(1):81–91.
24. Chang M, Canseco JA, Nicholson KJ, Patel N, Vaccaro AR. The role of machine learning in spine surgery: the future is now. *Front Surg*. 2020;7:54.
25. DeSole EM, Keck WL, Patel AA. The state of machine learning in spine surgery: a systematic review. *Clin Spine Surg*. 2022;35(2):80–9.
26. Han SS, Azad TD, Suarez PA, Ratliff JK. A machine learning approach for predictive models of adverse events following spine surgery. *Spine J*. 2019;19(11):1772–81.
27. Karhade AV, Cha TD, Fogel HA, Hershman SH, Tobert DG, Schoenfeld AJ, et al. Predicting prolonged opioid prescriptions in opioid-naïve lumbar spine surgery patients. *Spine J*. 2020;20(6):888–95.
28. Malik AT, Khan SN. Predictive modeling in spine surgery. *Ann Transl Med*. 2019;7(Suppl 5):S173.
29. Pedersen CF, Andersen MO, Carreon LY, Eiskjaer S. Applied machine learning for spine surgeons: predicting outcome for patients undergoing treatment for lumbar disc herniation using PRO data. *Global Spine J*. 2022;12(5):866–76.
30. Wilson JRF, Badhiwala JH, Moghaddamjou A, Martin AR, Fehlings MG. Degenerative cervical myelopathy; a review of the latest advances and future directions in management. *Neurospine*. 2019;16(3):494–505.
31. Yang SH, Kim CH, Lee CH, Ko YS, Won Y, Chung CK. C7 fracture as a complication of C7 dome-like laminectomy: impact on clinical and radiological outcomes and evaluation of the risk factors. *J Korean Neurosurg Soc*. 2021;64(4):575–84.
32. Kim DH, Lee CH, Ko YS, Yang SH, Kim CH, Park SB, Chung CK. The clinical implications and complications of anterior versus posterior surgery for multi-level cervical ossification of the posterior longitudinal ligament; an updated systematic review and meta-analysis. *Neurospine*. 2019;16(3):530–41.
33. Kim CH, Chung CK, Lee U, Choi Y, Park SB, Jung JM, et al. Postoperative changes in moderate to severe nonspecific low back pain after cervical myelopathy surgery. *World Neurosurg*. 2018;116:e429–35.
34. Yonenobu K, Abumi K, Nagata K, Taketomi E, Ueyama K. Interobserver and intraobserver reliability of the Japanese orthopaedic association scoring system for evaluation of cervical compression myelopathy. *Spine (Phila Pa 1976)*. 2001;26(17):1890–4. discussion 5.
35. Goh GS, Soh RCC, Yue WM, Guo CM, Tan SB, Chen JL. Determination of the patient acceptable symptom state for the Japanese Orthopaedic Association score in patients undergoing anterior cervical discectomy and fusion for cervical spondylotic myelopathy. *Spine J*. 2020;20(11):1785–94.
36. Mumtaz SL, Shamayleh A, Alshraideh H, Guella A. Improvement of dialysis dosing using big data analytics. *Healthc Inf Res*. 2023;29(2):174–85.
37. Feng J, Xu H, Mannor S, Yan S. Robust logistic regression and classification. *Adv Neural Inf Process Syst*. 2014;27.
38. Arayeshgari M, Najafi-Ghobadi S, Tarhsaz H, Parami S, Tapak L. Machine learning-based classifiers for the prediction of low birth weight. *Healthc Inf Res*. 2023;29(1):54–63.
39. Fung G, Mangasarian O, Shavlik J. Knowledge-based support vector machine classifiers. *Adv Neural Inf Process Syst*. 2002;15.
40. Zhang Z. Introduction to machine learning: k-nearest neighbors. *Ann Transl Med*. 2016;4:11.
41. Breiman L. Random forests. *Mach Learn*. 2001;45:5–32.
42. Chen T, He T, Benesty M, Khotilovich V, Tang Y, Cho H, et al. Xgboost: extreme gradient boosting. *R Package Version 04 – 2*. 2015;1(4):1–4.
43. Ramchoun H, Ghanou Y, Ettaouil M, Janati Idrissi MA. Multilayer perceptron: architecture optimization and training. *Int J Interact Multimed*. 2016;4:26–30.
44. Gardner MW, Dorling S. Artificial neural networks (the multilayer perceptron)—a review of applications in the atmospheric sciences. *Atmos Environ*. 1998;32(14–15):2627–36.
45. Medsker LR, Jain L. Recurrent neural networks. *Des Appl*. 2001;5:64–7.
46. Hochreiter S, Schmidhuber J. Long short-term memory. *Neural Comput*. 1997;9(8):1735–80.
47. Steeves JA, Tudor-Locke C, Murphy RA, King GA, Fitzhugh EC, Harris TB. Classification of occupational activity categories using accelerometry: NHANES 2003–2004. *Int J Behav Nutr Phys Act*. 2015;12:89.
48. Charlson ME, Pompei P, Ales KL, et al. A new method of classifying prognostic comorbidity in longitudinal studies: development and validation. *J Chronic Dis*. 1987;40(5):373–83.
49. Odukoya O, Nwaneri S, Odeniyi I, Akodu B, Oluwale E, Olorunfemi G, et al. Development and comparison of three data models for predicting diabetes mellitus using risk factors in a Nigerian population. *Healthc Inf Res*. 2022;28(1):58–67.
50. Lundberg SM, Lee S-I. A unified approach to interpreting model predictions. *Adv Neural Inf Process Syst*. 2017;30.
51. Xie X, Liu G, Cai Q, Wei P, Qu H. Multi-source sequential knowledge regression by using transfer RNN units. *Neural Netw*. 2019;119:151–61.
52. Wang K, Zhang K, Liu B, Chen W, Han M. Early prediction of sudden cardiac death risk with nested LSTM based on electrocardiogram sequential features. *BMC Med Inf Decis Mak*. 2024;24(1):94.
53. Daggubati LC, Eichberg DG, Ivan ME, Hanft S, Mansouri A, Komotar RJ, et al. Telemedicine for Outpatient Neurosurgical Oncology Care: lessons learned for the future during the COVID-19 pandemic. *World Neurosurg*. 2020;139:e859–63.
54. Eron L. Telemedicine: the future of outpatient therapy? *Clin Infect Dis*. 2010;51(Suppl 2):S224–30.
55. Goedeke J, Ertl A, Zoller D, Rohleder S, Muensterer OJ. Telemedicine for pediatric surgical outpatient follow-up: a prospective, randomized single-center trial. *J Pediatr Surg*. 2019;54(1):200.
56. Donnellan F, Hussain T, Aftab AR, McGurk C. Reducing unnecessary outpatient attendances. *Int J Health Care Qual Assur*. 2010;23(5):527–31.
57. Jamjanya S, Ruengorn C, Noppakun K, Thavorn K, Hutton B, Sood MM, et al. Temporal and external validation of the multidimensional scale-uremic pruritus in dialysis patients (UP-dial): a psychometric evaluation. *J Eur Acad Dermatol Venereol*. 2024;38(8):e694–7.
58. Zeng J, Zhang D, Lin S, Su X, Wang P, Zhao Y, et al. Comparative analysis of machine learning vs. traditional modeling approaches for predicting in-hospital mortality after cardiac surgery: temporal and spatial external validation based on a nationwide cardiac surgery registry. *Eur Heart J Qual Care Clin Outcomes*. 2024;10(2):121–31.
59. Martinez-Zayas G, Almeida FA, Yarmus L, Steinfert D, Lazarus DR, Simoff MJ, et al. Predicting lymph node metastasis in non-small cell lung cancer: prospective external and temporal validation of the HAL and HOMER models. *Chest*. 2021;160(3):1108–20.

Publisher's note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.